

Three-body calculation of the rate of reaction $p + p + e \rightarrow d + \nu_e$ in the Sun

B.F. Irgaziev,^{1,*} V.B. Belyaev,² and Jameel-Un Nabi¹

¹*GIK Institute of Engineering Sciences and Technology, Topi, Pakistan*

²*Joint Institute for Nuclear Research, Dubna, Russia*

(Dated: January 7, 2013)

Using expansion of the three-body wave function of the pep system in the initial state on hyperharmonic functions the rate of the $p + p + e^- \rightarrow d + \nu_e$ reaction in the Sun is calculated. The results of calculation of the flux at 1 AU are compared with the results of measurement made by the Borexino collaboration and Bahcall *et al.* theoretical predictions.

PACS numbers: 21.30.-x, 21.45.-v, 26.20.Cd, 26.65.+t, 27.10.+h, 96.60.Jw

I. INTRODUCTION

The study of the neutrino from the Sun is a multi-purpose problem. Indeed, the measurement of the solar neutrino flux incident on the Earth helps to clarify the properties of the neutrino, for example, the phenomenon of the oscillations and to determine its parameters (the mixing angle and the eigenmass of neutrino). Then, the value of the flux and its spectral properties contain in some cases information on the nuclear reactions that in the visible future cannot be observed in the laboratory. We mean here the reactions:

$$p + p \rightarrow d + e^+ + \nu_e \quad (1)$$

and

$$p + p + e^- \rightarrow d + \nu_e \quad (2)$$

Moreover, as emphasized by John N. Bahcall in his book and in a series the papers [1–5] (see also [6]) solar neutrinos bring the information about processes in the center of star connected with solar structure models.

Although the reaction (2), called the pep reaction, plays no significant role in hydrogen burning of stars, it is essential in detecting monoenergetic neutrinos of $E_\nu = 1.442$ MeV. Also, a measurement of the neutrino flux from reaction (2) can be useful for the determination of the parameters of the Standard Solar Model.

All above considerations contain sets of parameters used for fitting observable data in the framework of different models like star models, nuclear reactions models and so on. An example of such a type is the two-body model [7] for reaction (2) which is essentially three-body [8].

The purpose of this work is to exclude the model elements as much as possible in the description of reaction (2) and treat the initial state as a purely three-body state. From a strict point of view, the modern treatment of any nuclear reaction inside the Debye-sphere is still the model in a sense of the absence of the dynamical consideration

(it is the six-body problem in the Sun interior condition) of particles inside the sphere. We hope, however, that the three-body instead of the two-body treatment of reaction (2) is a step in the right direction.

Below, we concentrate ourselves mainly on the process (2) for the following reasons. Firstly, in 2012 the results of the first experimental observation of process (2) was announced [9], after more than 50 years of studies of solar neutrino problems. The second reason is connected with the absence of the three-body treatment of the initial state in the process (2).

There is a question concerning sensitivity to the choice of the nucleon-nucleon potential and related to these potentials wave functions of the bound state of the deuteron and the continuum state of the pep system.

With all this in mind we present below the treatment which takes into account all peculiarities of the pep three-body system. In Section II we start from the inputs for considered problem: the weak Hamiltonian and NN -potentials, then in Section III, we consider solution of the Schrödinger equation to determine the pep wave function of the initial state. In Section IV, we present the calculation of the probability of the pep reaction and the astrophysical S_{pep} factor taking into account the Coulomb and strong interactions simultaneously. The results for the rate of the process and fluxes of the neutrino are discussed in Section V and conclusions are presented in Section VI.

II. INPUTS

The electron capture by the nuclear system can be described by the following non-relativistic effective weak Hamiltonian [10]

$$H_w = \frac{1}{\sqrt{2}} \tau^{(+)} \frac{1 - \boldsymbol{\sigma} \cdot \boldsymbol{\nu}_1}{\sqrt{2}} \sum_{i=1}^A \tau_i^{(-)} [G_V \mathbf{1} \cdot \mathbf{1}_i + G_A \boldsymbol{\sigma} \cdot \boldsymbol{\sigma}_i - G_P \boldsymbol{\sigma} \cdot \boldsymbol{\nu}_1 \boldsymbol{\sigma}_i \cdot \boldsymbol{\nu}_1] \delta(\mathbf{r} - \mathbf{r}_i), \quad (3)$$

where $\boldsymbol{\nu}_1 = \boldsymbol{\nu}/\nu$ (ν is the neutrino momentum); $\mathbf{1}, \mathbf{1}_i$, $\boldsymbol{\sigma}$ and $\boldsymbol{\sigma}_i$ are the 2×2 matrix unit operators and spin angular momentum operators for the lepton and i th nucleon; $\mathbf{r}$ and $\mathbf{r}_i$ are the space coordinates of the lepton and i th

*Electronic address: irgaziev@yahoo.com

nucleon; $\tau_i^{(+)}$, $\tau_i^{(-)}$ are the isobaric-spin operators which transform a lepton electron state into a lepton neutrino state and i th nucleon proton state into the i th nucleon neutron state; G_V , G_A and G_P are the vector, axial vector and “induced” pseudoscalar coupling constants, respectively. We take $G_V/(\hbar c)^3 = 1.153 \times 10^{-11} \text{ GeV}^{-2}$ and $G_A/(\hbar c)^3 = -1.454 \times 10^{-11} \text{ GeV}^{-2}$ [11]. We can simplify the weak Hamiltonian for the pep system: the last term in Eq. (3) can be neglected because the emitted neutrino has energy $E_\nu = 1.442 \text{ MeV}$ and this term encloses factor $\nu/2m_p$, where m_p is the proton mass; the $pep \rightarrow d + \nu_e$ transition satisfies the Gamow-Teller selection rule; therefore, the first term of Eq. (3) does not give contribution to the matrix element of transition; and at last we take into account that the electron neutrino has spin opposite to its momentum ν . Finally, for the weak Hamiltonian we get

$$H_w = \tau^{(+)} G_A \sum_{i=1}^A \tau_i^{(-)} \boldsymbol{\sigma} \cdot \boldsymbol{\sigma}_i \delta(\mathbf{r} - \mathbf{r}_i). \quad (4)$$

With this Hamiltonian of Eq. (4) we obtain the electron capture transition matrix element.

The energy of thermalization of particles in the interior of the Sun corresponds to $E \sim 1.3 \text{ keV}$ which is small on the nuclear energy scale. Therefore, we can use the simple NN -potential which describes correctly the low energy data of the nucleon-nucleon system. We apply the Gauss and the Yukawa potentials [12]. We fit the parameters of these potentials to get the values of the deuteron energy, the scattering lengths and the effective ranges for pp and pn scattering.

For the Gauss potential

$$V(r) = -V_0 \exp(-r^2/R_N^2). \quad (5)$$

the calculation with the fitted parameters

$$V_0^s = 30.36 \text{ MeV}, \quad R_N^s = 1.816 \text{ fm}$$

gives the scattering length $^s a_{pp} = -7.884 \text{ fm}$ and the effective range $^s r_{pp} = 2.678 \text{ fm}$ for pp scattering at the singlet state (s), while the parameters

$$V_0^t = 60.572 \text{ MeV}, \quad R_N^t = 1.65 \text{ fm},$$

lead to the scattering length $^t a_{pp} = 5.484 \text{ fm}$, the effective range $^t r_{pp} = 1.85 \text{ fm}$ for pp scattering at the triplet state (t) and the binding deuteron energy $\varepsilon_d = 2.225 \text{ MeV}$.

The second potential is the Yukawa potential

$$V(r) = -\frac{V_0}{r/R_N} \exp(-r/R_N) \quad (6)$$

with the parameters for the singlet state:

$$V_0^s = 44.05 \text{ MeV}, \quad R_N^s = 1.206 \text{ fm},$$

and for the triplet state:

$$V_0^t = 53.27 \text{ MeV}, \quad R_N^t = 2.43 \text{ fm}.$$

The results of calculation of the scattering lengths and effective ranges are:

$$\begin{aligned} ^s a_{pp} &= -7.782 \text{ fm}, & ^s r_{pp} &= 2.868 \text{ fm}, \\ ^t a_{np} &= 5.626 \text{ fm}, & ^t r_{np} &= 1.895 \text{ fm}. \end{aligned} \quad (7)$$

To find the flux of neutrino we must use some Standard Solar Model (SSM). There are several SSM which are in good agreement with the helioseismologically determined sound speed, temperature and density of elements as a function of solar radius, the depth of the convective zone, the surface helium abundance, and so on. We applied data of parameters presented in the model BS05(OP) [13]. The results of Bahcall *et al.* [14] shows that the flux from the pep reaction is not sensitive to the type of SSM.

III. THE WAVE FUNCTION OF THE pep INITIAL STATE

We note that Bahcall and May [7] used the pep wave function in a factorized form as the product of the wave function of the relative motion of two protons and the wave function of the electron moving in the Coulomb field of these protons. However, such a representation is not a quite correct procedure due to the long-range nature of the Coulomb interaction, even for the asymptotic behavior of the wave function when electron is at a large distance from the protons [15, 16]. All the more, we cannot perform such a factorization at a small relative distance where we need to know the wave function with a sufficient accuracy to get a good accuracy of the calculation for the transition matrix element of the process $pep \rightarrow d\nu_e$. Also, there is a problem with the total angular momentum of pep , because the moments of subsystems are not conserved. Fortunately, the relative angular moment of two nucleons in the initial and final states is zero as angular moments of the electron and the neutrino; therefore, the last problem did not arise in the Bahcall calculations of the pep reaction.

We use the hyperspherical harmonics [17, 18] for expansion of the pep wave function in the initial state and solve directly the three-body Schrödinger equation and, therefore, we are free of the problems which we mentioned above. As in the nonrelativistic approach the orbital and spin moments are conserved independently of one another, we can expand the spatial part of the pep three-particle wave function over hyperspherical functions and we obtain the linked system of the radial differential equations. Derivation of the system radial equations is given in Appendix A. To proceed further, we now use the system of Eqs (A14) for the radial wave functions. Since the total energy of the pep system is low, the main contribution to the three-particle wave function gives the states with the zero relative orbital moments. Owing to the centrifugal potential and $\kappa\rho \ll 1$ at small distance where we need to calculate the wave function with higher

accuracy, contributions of the components with the hypermomenta $K > 0$ should be suppressed in the total wave function. Taking into account these conditions, we need to find the solution of the single equation only for $K = l_x = l_y = 0$ with omitting the non-diagonal terms. We omit all the indices because they correspond to zero values of the quantum numbers and get the equation for the radial wave function $U(\rho)$:

$$\frac{d^2 U(\rho)}{d\rho^2} + \frac{1}{\rho} \frac{dU}{d\rho} - \left(\mathcal{V}(\rho) + \frac{4}{\rho^2} - \kappa^2 \right) U(\rho) = 0, \quad (8)$$

where $\kappa^2 = 2\mu_{23}E > 0$ (E is the total energy of the pep system);

$$\mathcal{V}(\rho) = \mathcal{V}^N(\rho) + \mathcal{V}^C(\rho), \quad (9)$$

$$\mathcal{V}^C(\rho) = \frac{32\mu_{23}}{3\pi\hbar^2} \frac{(a_1 + a_2 + a_3)}{\rho} \equiv \frac{2\eta_3\kappa}{\rho}. \quad (10)$$

Here η_3 is the three-body Coulomb parameter which is defined as

$$\eta_3 = \frac{16\mu_{23}}{3\pi\hbar^2\kappa} (a_1 + a_2 + a_3), \quad (11)$$

$$a_1 = \sqrt{\frac{m_2 m_3}{\mu_{23}(m_2 + m_3)}} e^2 \equiv e^2, \quad (12)$$

$$a_2 = -\sqrt{\frac{m_1 m_3}{\mu_{23}(m_1 + m_3)}} e^2 \simeq -\sqrt{\frac{2m_1}{m_3}} e^2, \quad (13)$$

$$a_3 = -\sqrt{\frac{m_1 m_2}{\mu_{23}(m_1 + m_2)}} e^2 \simeq -\sqrt{\frac{2m_1}{m_2}} e^2. \quad (14)$$

The matrix element of the nuclear potential $\mathcal{V}^N$ is the following: for the Gauss potential

$$\mathcal{V}^N(\rho) = -\frac{8\mu_{23}V_0}{\hbar^2} \frac{\exp\left(-\frac{\rho^2}{2R_N^2}\right) I_1\left(\frac{\rho^2}{2R_N^2}\right)}{\rho^2/R_N^2}, \quad (15)$$

and for the Yukawa potential

$$\mathcal{V}^N(\rho) = -\frac{16\mu_{23}V_0}{\hbar^2} \frac{\frac{2\rho}{3\pi R_N} - I_2\left(\frac{\rho}{R_N}\right) + L_2\left(\frac{\rho}{R_N}\right)}{\rho^2/R_N^2}, \quad (16)$$

where $I_n(z)$ is the modified Bessel function of the first kind, and $L_n(z)$ is the modified Struve function.

To find a unique solution of Eq. (8) for the continuous state of the pep system, we must determine boundary conditions. Instead, to define the function $U(\rho)$ and its derivative at the origin ($\rho = 0$), we define the wave function in a point ρ_0 close to zero because we know behaviour of the wave function near the origin. We can take boundary conditions as

$$U(\rho_0) = J_2(\kappa_0\rho_0), \quad U'(\rho_0) = \kappa_0 J_2'(\kappa_0\rho_0), \quad (17)$$

where $\kappa_0 = \sqrt{\kappa^2 + |\mathcal{V}_N(\rho_0)|}$.

The wave function $U(\rho)$ at large distance, where the nuclear interaction is negligible, has the following asymptotics:

$$U(\rho) \xrightarrow{\rho \rightarrow \infty} e^{i\delta_3} \cos \delta_3 (F_{00}(\kappa\rho) - \tan \delta_3 G_{00}(\kappa\rho)), \quad (18)$$




FIG. 1: (Color online) The pep radial wave function (solid curve) obtained by solving Eq. (8) and its asymptotics at $\rho \gg R_N$ (dashed curve). The pure Coulomb wave is dot-dashed curve. $E_{pep} = 6$ keV.

where δ_3 is the three-body scattering phase shift, $F_{00}(\kappa\rho)$ ($G_{00}(\kappa\rho)$) is a three-body regular (irregular) Coulomb wave function. The Coulomb wave functions have asymptotics [17] at the large distance

$$F_{00}(z) = \sqrt{\frac{2}{\pi z}} \cos\left(z - \eta_3 \ln 2z - \frac{5}{4}\pi + \sigma^C\right), \quad (19)$$

$$G_{00}(z) = \sqrt{\frac{2}{\pi z}} \sin\left(z - \eta_3 \ln 2z - \frac{5}{4}\pi + \sigma^C\right), \quad (20)$$

where $z = \kappa\rho$ and $\sigma^C = \arg \Gamma(5/2 + i\eta_3)$ is the three-body Coulomb scattering phase shift when hypermoment K and the total angular momentum L are equal to zero, and $\Gamma(z)$ is the gamma function.

We solve Eq. (8) using the boundary conditions (17) and then matching the logarithmic derivative of the solution in the asymptotic region with the logarithmic derivative of the asymptotic solution (Eq. (18)) we define the three-body phase shift δ_3 which depends on the total energy E of the pep system. Of course, this phase shift is very small because $\kappa\rho_N \ll 1$, where ρ_N is equal to the radius of NN interaction. Matching the condition of the numerically obtained solution of the Schrödinger equation with its asymptotics also allows us to find the normalization factor. At great distances the ratio of the unnormalized solution to the asymptotic function becomes a constant.

Figure 1 shows the results of calculation of the pep radial wave function (solid curve) at $E = 6$ keV and its asymptotics for the Gauss NN potential (dashed curve). Since the value of the total energy of the system is small, the scattering phase will be small too. Therefore the asymptotics of the solution can be described by a pure three-body Coulomb wave (dot-dashed curve). We obtain the same results for the Yukawa NN potential. We need to know the three-body wave function with high accuracy in the interval $0 < \rho < 35$ fm because the deuteron

wave function cuts integration at the distance 35 fm when we calculate the matrix element of the $pep \rightarrow d + \nu$ transition.

IV. THE PROBABILITY OF THE $pep \rightarrow d + \nu$ REACTION AND ASTROPHYSICAL S FACTOR

Using the weak Hamiltonian (Eq. (4)) we can write the matrix element of the pep reaction as

$$H_{if} = G_A < \varphi_\nu | \sigma \tau^{(+)} | \varphi_e > \times \sum_{i=1}^2 < \Psi_d | \sigma_i \tau_i^{(-)} | \Psi_{pep} >, \quad (21)$$

where φ_e and φ_ν are the spin functions of the electron and neutrino, respectively; Ψ_d is the wave function of the deuteron, while Ψ_{pep} is the three-body wave function of the pep system in a continuous state.

Since the weak interaction is small, the first-order perturbation theory can be applied for calculation of the probability of reaction per unit of time P_3 ; therefore, we have

$$P_3 = \frac{2\pi}{\hbar} \overline{|H_w|^2} \rho(E_\nu), \quad (22)$$

where the overline means summation over spins in the final state and averaging over spins in the initial state, and the density of neutrino states is

$$\rho(E_\nu) = \frac{E_\nu^2}{2\pi^2 \hbar^3 c^3}. \quad (23)$$

Here E_ν , c are the neutrino energy and light speed, respectively. Using the energy conservation law we define

$$E_\nu = E_{pp} + E_e - (M_d - 2m_p - m_e)c^2,$$

where E_{pp} is the kinetic energy of proton-proton relative motion, M_d , m_p , m_e are the deuteron, proton and electron masses, respectively. If we put kinetic energies of the particles to zero, we get $E_\nu = 1.442$ MeV.

Making averaging in Eq. (22) and using the Jacobi coordinates for the particles defined by Eq. (A5) of Appendix A we obtain

$$P_3 = \frac{3E_\nu^2 G_A^2}{\pi \hbar^4 c^3} \left| \int \Psi_d^*(\mathbf{x}_1) \Psi_{pep}(\mathbf{x}_1, \mathbf{y}_1) |_{y_1=y_{10}} d^3 x_1 \right|^2. \quad (24)$$

Note, the presence of the delta-function in the weak Hamiltonian (4) leads to calculation of the integral over variable $\mathbf{y}_1$ to the significance of the pep wave function at the distance $y_{10} = \sqrt{\frac{m_1(m_2+m_3)}{\mu_{23}(m_1+m_2+m_3)}} x_1/2 \simeq \sqrt{\frac{m_e}{2m_p}} x_1$.

The deuteron wave function can be written in the form

$$\Psi_d(\mathbf{x}_1) = \frac{\chi_d(x_1)}{x_1} Y_{00}(\mathbf{x}_1), \quad (25)$$

where the radial wave function must have a correct value of the asymptotic normalization coefficient to get a correct value of the probability of the reaction. Using Eq. (25) and taking into account only the first term in the expansion of the pep wave function over the hyperharmonics functions we obtain for the overlap integral

$$\int \Psi_d^*(\mathbf{x}_1) \Psi_{pep}(\mathbf{x}_1, \mathbf{y}_1) |_{y_1=y_{10}} = \frac{16\sqrt{\pi}}{\kappa} e^{i\delta_3} \cos \delta_3 \int_0^\infty \chi_d(x_1) U(x_1) dx_1. \quad (26)$$

Here the three-body radial wave function is calculated at the point $\rho = \sqrt{x_1^2 + y_{10}^2} \simeq x_1$, and δ_3 is the three-body nuclear scattering phase shift modified by the Coulomb interactions.

Usually, in the nuclear astrophysics for parameterizations of a two-body reaction cross section the astrophysical S factor is used [19]. To find the S factor from the cross section at a small energy region, the Gamow factor is extracted from the cross section, i.e.

$$\sigma(E) = \frac{S(E)}{E} e^{-2\pi\eta}, \quad (27)$$

where $\eta = Z_1 Z_2 e^2 / (\hbar v)$ is the Sommerfeld (Coulomb) factor for the colliding nuclei with charges Z_1 and Z_2 having the relative velocity v . A similar procedure can be carried out for the reaction with three particles in the initial state because the three-body radial wave function $U_{KL}^{l_x l_y}(\rho)$ in the continuum state at a distance larger than nuclear interaction radius encloses factor $\exp(-\pi\eta_3/2) | \Gamma(K + 5/2 + i\eta_3) |$. Therefore we define the astrophysical S_{pep} factor for the pep reaction as

$$P_3(E) = G_0(E) S_{pep}(E), \quad (28)$$

$$G_0(E) = \frac{2\pi e^{-2\pi\eta_3} (\frac{1}{4} + \eta_3^2) (\frac{9}{4} + \eta_3^2)}{1 + e^{-2\pi\eta_3}}. \quad (29)$$

$S_{pep}(E)$ is almost a linearly varying function of E and $S_{pep}(0)$ is not equal to zero, like the astrophysical S_{pp} factor for two-body reactions. Note, the unit of S_{pep} coincides with the unit of P_3 because $G_0(E)$ is dimensionless and it is the Gamow factor for the pep reaction. If we use expansion of $S_{pep}(E)$ over E as

$$S_{pep}(E) = S_0 + S_1 E + S_2 E^2, \quad (30)$$

we obtain the following results for the value of the coefficients:

For the Gauss potentials

$$\begin{aligned} S_0 &= 2.38 \times 10^{10} \text{ fm}^6/\text{s}; \\ S_1 &= 3.03 \times 10^{10} \text{ fm}^6/(\text{MeVs}); \\ S_2 &= 1.45 \times 10^{10} \text{ fm}^6/(\text{MeV}^2\text{s}); \end{aligned} \quad (31)$$

and for the Yukawa potentials

$$\begin{aligned} S_0 &= 2.33 \times 10^{10} \text{ fm}^6/\text{s}; \\ S_1 &= 3.01 \times 10^{10} \text{ fm}^6/(\text{MeVs}); \\ S_2 &= 1.78 \times 10^{10} \text{ fm}^6/(\text{MeV}^2\text{s}). \end{aligned} \quad (32)$$




FIG. 2: (Color online) The pep astrophysical S_{pep} factor as a function of energy calculated by the Gauss (solid line) and Yukawa (dashed line) NN potentials.

Behavior of the S_{pep} astrophysical factor on energy is presented in Fig. 2 where linear dependence of the S factor is seen clearly for both potentials. The small difference between the S_{pep} factors for the Gauss and Yukawa potentials can be explained by small differences of the obtained low energy scattering parameters. To check the validity of the results of our calculations of the pep reaction, we calculated the astrophysical S_{pp} factor for the pp reaction, according to the conventional definition (Eq. (27)) by the same NN potentials. If S_{pp} is approximated by a polynomial

$$S_{pp}(E) = S_0 + S_1 E + S_2 E^2, \quad (33)$$

we get the following results:

For the Gauss potentials

$$\begin{aligned} S_0 &= 4.21 \times 10^{-25} \text{ MeVb}; \\ S_1 &= 4.75 \times 10^{-24} \text{ b}; \\ S_2 &= 3.02 \times 10^{-23} \text{ MeV}^{-1} \text{ b}; \end{aligned} \quad (34)$$

and for the Yukawa potentials

$$\begin{aligned} S_0 &= 4.11 \times 10^{-25} \text{ MeVb}; \\ S_1 &= 4.64 \times 10^{-24} \text{ b}; \\ S_2 &= 2.93 \times 10^{-23} \text{ MeV}^{-1} \text{ b}. \end{aligned} \quad (35)$$

These results are very close to data presented in [19].

V. RATE OF THE pep REACTION AND THE SOLAR NEUTRINO FLUX

The number of the pep reactions per unit of time per unit of the solar interior volumes of protons and electrons

(a rate constant) is

$$\mathcal{K}_{pep} = \langle P_3 \rangle = \int_0^\infty \int_0^\infty \int_0^\infty \varphi(v_e) \varphi(v_{p_1}) \varphi(v_{p_2}) P_3(E) dv_e dv_{p_1} dv_{p_2}, \quad (36)$$

where $\varphi(v) = 4\pi v^2 \left(\frac{m}{2\pi kT}\right)^{3/2} \exp\left(-\frac{mv^2}{2kT}\right)$ is the Maxwell-Boltzmann distribution function for particles of mass m and velocity v and k is the Boltzmann constant. Excluding the velocity of the center of mass of the pep system by transferring to the relative velocities and the fact that the center of mass of the pep system is almost the same as the center of pp we get

$$\mathcal{K}_{pep} = \int_0^\infty \int_0^\infty \varphi(v_e) \varphi(v_{pp}) P_3(E) dv_e dv_{pp}, \quad (37)$$

where $\varphi(v_{pp}) = 4\pi v_{pp}^2 \left(\frac{\mu_{pp}}{2\pi kT}\right)^{3/2} \exp\left(-\frac{\mu_{pp} v_{pp}^2}{2kT}\right)$, v_{pp} is the relative velocity of protons and $\mu_{pp} = m_p/2$. Taking into account that

$$E = \frac{\mu_{pp} v_{pp}^2}{2} + \frac{m_e v_e^2}{2}$$

and making transformations

$$v_{pp} = V \cos \alpha, \quad v_e = \sqrt{\frac{\mu_{pp}}{m_e}} V \sin \alpha, \quad E = \frac{\mu_{pp} V^2}{2}$$

we obtain

$$\mathcal{K}_{pep} = \frac{1}{(kT)^3} \int_0^\infty e^{-E/kT} P_3(E) E^2 dE. \quad (38)$$

Using Eqs. (28), (29) we finally obtain

$$\mathcal{K}_{pep} = \sqrt{\frac{8}{\pi \mu_{pp}}} \frac{c}{(kT)^{3/2}} \int_0^\infty e^{-2\pi\eta_3 - E/kT} S_{pep}(E) E^2 dE. \quad (39)$$

We note that the rate constant depends on the temperature, the nature of the reactants, but does not depend on their concentration. If we define the Gamow energy as

$$E_G = \frac{1}{2} \left(\frac{2\pi e^2 z_{eff}}{\hbar} \right)^2, \quad (40)$$

the integrand of Eq. (39) has the maximum at the energy

$$E_{max} = \left(\frac{1}{2} kT \sqrt{E_G} \right)^{2/3}. \quad (41)$$

Here z_{eff} is defined through Eqs. (12), (13) and (14) and it equals to

$$z_{eff} = \frac{16}{3\pi} (a_1 + a_2 + a_3). \quad (42)$$




FIG. 3: (Color online) The *pep* (solid line) and *pp* (dashed line) rates versus the radius of the interior of the Sun. The results are presented for the Gauss *NN* potentials. The behaviour of the rate curves for the Yukawa *NN* potentials is very close to the displayed curves. The results are presented in units of 10^7 (*pp*) and 10^5 (*pep*) $\text{cm}^{-3}\text{s}^{-1}$. $R_{\odot}$ is the radius of the Sun.

Note, if we take $z_{eff} = 1$ we obtain the point of maximum of the integrand in the equation of the *pp* reaction rate constant.

To obtain the rate of reactions, the rate constant must be multiplied by the density of the reactants,

$$\mathcal{R}_{pep} = \mathcal{K}_{pep} n_p^2 n_e, \quad (43)$$

where n_e and n_p are the numbers of electrons and protons in the unit of volume.

All solar parameters like temperature, densities of protons and electrons vary with the radius of interior of the Sun. The results for the neutrino fluxes presented by Bahcall *et al.* [5] show that the fluxes from the *pp* and *pep* reaction are not sensitive to the considered solar models. Therefore, for calculation of the rates of the *pp* and *pep* reaction we apply the BS05(OP) model [6].

Figure 3 shows the rate of the *pp* and *pep* reaction as a function of the solar interior radius for the Gauss *NN* potentials. We obtain the same behaviour in the case of the Yukawa *NN* potentials. We see that the rate of the *pep* reactions is more than a hundred times less than the rate of the *pp* reaction for the whole distance from the center of the Sun, and reactions occur in the core of the Sun where the temperature and density are higher.

Integrating the reaction rate (43) over the volume of the Sun we find the total flux of neutrinos emitted by the Sun. Dividing this total flux by the area of the sphere of the radius of one astronomical unit (AU) we obtain the neutrino flux Φ passing through a unit area of the Earth surface. The results of calculations of the neutrino fluxes are presented in Table I. Note, the results for the neutrino flux from the *pp* reaction are close to the results obtained by Bahcall *et al.* while there are differences between the fluxes from the *pep* reactions.

TABLE I: Predicted fluxes Φ_{pp} and Φ_{pep} (without survival probability), in units of 10^{10} (*pp*), 10^8 (*pep*) $\text{cm}^{-2}\text{s}^{-1}$.

Standard Solar Model	Φ_{pp}	Φ_{pep}	Φ_{pp}/Φ_{pep}	References
BS05(OP)	6.20	2.04	304	our results with Gauss potential
BS05(OP)	6.05	1.99	304	our results with Yukawa potential
BP04(Yale)	5.94	1.40	424	[4]
BP04(Garching)	5.94	1.41	421	[4]
BS04	5.94	1.40	424	[4]
BS05(14N)	5.99	1.42	421	[4]
BS05(OP)	5.99	1.42	421	[4]
BS05(AGS,OP)	6.06	1.45	418	[4]
BS05(AGS,OPAL)	6.05	1.45	417	[4]

The Borexino collaboration announced the results of the neutrino flux measurement: $\Phi_{pep} = (1.6 \pm 0.3) \times 10^8 \text{cm}^{-2}\text{s}^{-1}$ [9]. Taking into account that the survival probability of the neutrino (due to the neutrino oscillation) in the *pep* reaction predicted by the Borexino collaboration equals $P = 0.62 \pm 0.17$ at 1.44 MeV [9], we find that the neutrino flux at 1 AU should be equal to $\Phi_{pep} = (1.27 \pm 0.35) \times 10^8 \text{cm}^{-2}\text{s}^{-1}$ for the Gauss potential, and $\Phi_{pep} = (1.24 \pm 0.34) \times 10^8 \text{cm}^{-2}\text{s}^{-1}$ for the Yukawa potential.

We note that our results for the neutrino fluxes from the *pep* reaction obtained by taking into account the survival probability lie within the confidence interval of the experimental data. At the same time, the Bahcall results are out of this limit at all fluxes listed in Table I. Comparing our calculated low-energy parameters for the Gauss and Yukawa potentials, we see that they differ by 2-7%, and the neutrino fluxes from the *pp* have 2% difference too, and the results of Φ_{pp} of Bahcall *et al.* shown in Table I for all SSM are not more than 2-4% from our calculated Φ_{pp} . Therefore, we may conclude that the dependence of the results on the type of *NN* potentials is very weak. However, the difference in the results obtained by us and Bahcall *et al.* can reach up to 39-45%, which means strong sensitivity to the choice of the wave functions of the initial three-body state of the *pep* system.

VI. CONCLUSION

In the framework of the three-body approach the probability of the process $pep \rightarrow d + \nu_e$ under the condition of the solar core has been found. The rate of the above process and neutrino flux are found and compared with the Borexino experiment and previous calculations. The

value of the neutrino flux obtained from the *pep* reaction in the three-body treatment appeared to be $\sim 40\%$ larger as compared to the Bahcall *et al.* value. This can be understood as a correct description of the movement of the electron producing the screening effect between protons. In this work we have introduced the astrophysical S factor for the three-body reaction which is an analog of the S factor introduced for binary processes. To discriminate between different star models on the basis of our results, it is necessary to essentially reduce experimental errors in the above experiment.

Acknowledgments

The authors are very grateful to Prof. O.A. Zaimidzora for fruitful discussions and useful comments.

Appendix A: Schrödinger equation

The Hamiltonian of the *pep* system is

$$H = H_0 + V_{23}^N + V_{123}^C, \quad (\text{A1})$$

where H_0 is kinetic energy operator:

$$H_0 = -\frac{\hbar^2}{2\mu_{23}}\Delta_{23} - \frac{\hbar^2}{2M_{1(23)}}\Delta_{1(23)}, \quad (\text{A2})$$

subscript 1 means electron, while 2 and 3 correspond to two protons; μ_{23} is the reduced mass of two protons, while $M_{1(23)}$ is the reduced mass of the system of two protons and electron;

$$V_{123}^C = V_{12}^C + V_{23}^C + V_{31}^C \quad (\text{A3})$$

is the sum of the Coulomb potentials in the *pep* system, and V_{23}^N is nuclear potential of interaction between two nucleons.

The wave function of the initial state of the *pep* system is the eigenfunction of the Hamiltonian (A1), which has a continuous spectrum of energy and satisfies the Schrödinger equation

$$H\Psi(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3) = E\Psi(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3). \quad (\text{A4})$$

Let us define the Jacobi coordinates:

$$\begin{aligned} \mathbf{x}_i &= \sqrt{\frac{m_j m_k}{(m_j + m_k)\mu_{23}}}(\mathbf{r}_j - \mathbf{r}_k) \\ \mathbf{y}_i &= \sqrt{\frac{m_i(m_j + m_k)}{(m_1 + m_2 + m_3)\mu_{23}}} \times \\ &\quad \left(-\mathbf{r}_i + \frac{m_j \mathbf{r}_j + m_k \mathbf{r}_k}{m_j + m_k}\right), \end{aligned} \quad (\text{A5})$$

where $\mu_{23} = m_2 m_3 / (m_2 + m_3)$ is the reduced mass of two protons, indices $ijk=123, 231, \text{ or } 312$ and $m_i(\mathbf{r}_i)$ is the mass (coordinate) of particle i . In introducing

coordinates the Hamiltonian H_0 of the free motion of particles can be written as

$$\begin{aligned} H_0 &= -\frac{\hbar^2}{\mu_{23}}(\Delta_{\mathbf{x}_1} + \Delta_{\mathbf{y}_1}) \equiv -\frac{\hbar^2}{\mu_{23}}(\Delta_{\mathbf{x}_2} + \Delta_{\mathbf{y}_2}) \\ &\equiv -\frac{\hbar^2}{\mu_{23}}(\Delta_{\mathbf{x}_3} + \Delta_{\mathbf{y}_3}). \end{aligned} \quad (\text{A6})$$

We define the square of the hyperradius and hyperangle as:

$$\begin{aligned} \rho^2 &= x_1^2 + y_1^2 \equiv x_2^2 + y_2^2 \equiv x_3^2 + y_3^2, \quad \rho \in [0, \infty), \\ x_i &= \rho \cos \alpha_i, \quad y_i = \rho \sin \alpha_i, \quad \alpha_i \in [0, \pi/2]. \end{aligned}$$

In the variables of hyperradius and a set of angles $\Omega_i = (\hat{\mathbf{x}}_i, \hat{\mathbf{y}}_i, \alpha_i)$ ($\hat{\mathbf{x}}_i$ and $\hat{\mathbf{y}}_i$ are unit vectors determining the azimuthal and polar angles), we can rewrite the operator H_0 as

$$H_0 = -\frac{\hbar^2}{2\mu_{23}}\left(\frac{\partial^2}{\partial \rho^2} + \frac{5}{\rho} \frac{\partial}{\partial \rho} - \frac{1}{\rho^2} K^2(\Omega_i)\right), \quad (\text{A7})$$

where the operator $K^2(\Omega_i)$ is

$$\begin{aligned} K^2(\Omega_i) &= -\frac{\partial^2}{\partial \alpha_i^2} - 4 \cot 2\alpha_i \frac{\partial}{\partial \alpha_i} + \\ &\quad \frac{1}{\cos^2 \alpha_i} l^2(\hat{\mathbf{x}}_i) + \frac{1}{\sin^2 \alpha_i} l^2(\hat{\mathbf{y}}_i), \end{aligned} \quad (\text{A8})$$

where $l^2 = -\Delta_{\theta, \varphi}$ is an angular part of the Laplace operator.

The hyperspherical function is defined as the solution of the equation [17]

$$K^2 \Phi(\Omega_i) = K(K+4) \Phi(\Omega_i), \quad K = 0, 1, 2, 3, \dots \quad (\text{A9})$$

The quantum number K is called hypermoment, the eigenfunction $\Phi(\Omega_i)$ is

$$\begin{aligned} \Phi(\Omega_i) &\equiv \Phi_{KLM}^{l_{x_i} l_{y_i}}(\Omega_i) \\ &= \sum_{m_{x_i} m_{y_i}} (l_{x_i} m_{x_i} l_{y_i} m_{y_i} | LM) \Phi_{KLM}^{l_{x_i} l_{y_i} m_{x_i} m_{y_i}}(\Omega_i), \end{aligned} \quad (\text{A10})$$

where the function $\Phi_{KLM}^{l_{x_i} l_{y_i} m_{x_i} m_{y_i}}(\Omega_i)$ is

$$\begin{aligned} \Phi_{KLM}^{l_{x_i} l_{y_i} m_{x_i} m_{y_i}}(\Omega_i) &= N_K^{l_{x_i} l_{y_i}} (\cos \alpha_i)^{l_{x_i}} (\sin \alpha_i)^{l_{y_i}} \times \\ &\quad P_n^{l_{y_i}+1/2, l_{x_i}+1/2}(\cos 2\alpha_i) Y_{l_{x_i} m_{x_i}}(\hat{\mathbf{x}}_i) Y_{l_{y_i} m_{y_i}}(\hat{\mathbf{y}}_i); \end{aligned} \quad (\text{A11})$$

$(l_{x_i} m_{x_i} l_{y_i} m_{y_i} | LM)$ is the Clebsch-Gordon coefficient; $Y_{l_x m_x}(\hat{\mathbf{x}})$ is the spherical function;

$$N_K^{l_{x_i} l_{y_i}} = \sqrt{\frac{2n!(K+2)(n+l_{x_i}+l_{y_i}+1)!}{\Gamma(n+l_{x_i}+3/2)\Gamma(n+l_{y_i}+3/2)}}, \quad (\text{A12})$$

$n = (1/2)(K - l_{x_i} - l_{y_i})$ must be an integer number; $P_n^{l_x l_y}$ is Jacobi polynomial.

Let $\mathbf{p}$ and $\mathbf{q}$ be conjugate momenta to the coordinates $\mathbf{x}$ and $\mathbf{y}$. Then determining square of the wave number $\kappa^2 = 2\mu_{23}E/\hbar^2$ (E is the total energy of the *pep* system in the c.m. frame) for the *pep* system in the continuous state, removing a free motion of the center of mass of the *pep* system from Eq. (A4) and using the following expansion of the *pep* wave function of a continuous spectrum over the hyperharmonics functions

$$\Psi_{\mathbf{p},\mathbf{q}}(\mathbf{x},\mathbf{y}) = (2\pi)^3 \sum_{Kl_xl_yLM} i^K \frac{U_{KL}^{l_xl_y}(\rho)}{(\kappa\rho)^2} \times \Phi_{KLM}^{l_xl_y}(\Omega_\rho) \Phi_{KLM}^{*l_xl_y}(\Omega_\kappa) \quad (\text{A13})$$

we get the radial Schrödinger equation:

$$\begin{aligned} \frac{d^2 U_{KL}^{l_xl_y}(\rho)}{d\rho^2} + \frac{1}{\rho} \frac{dU_{KL}^{l_xl_y}(\rho)}{d\rho} - \left[\frac{(K+2)^2}{\rho^2} - \kappa^2 \right] U_{KL}^{l_xl_y}(\rho) \\ = \sum_{K'l'_x l'_y} \mathcal{V}_{KK'LM}^{l_xl_y;l'_x l'_y}(\rho) U_{K'L}^{l'_x l'_y}(\rho), \end{aligned} \quad (\text{A14})$$

where superscript (*) denotes taking complex conjugate

of the function, and $\Phi_{KLM}^{l_xl_y}(\Omega_\rho) \equiv \Phi_{KLM}^{l_xl_y}(\Omega_1)$, $\Phi_{KLM}^{l_xl_y}(\Omega_\kappa)$ is defined on a hypersphere of unit radius in a six dimensional momentum space. The matrix element $\mathcal{V}_{KK'LM}^{l_xl_y;l'_x l'_y}(\rho)$ equals to

$$\begin{aligned} \mathcal{V}_{KK'LM}^{l_xl_y;l'_x l'_y}(\rho) = \frac{2\mu_{23}}{\hbar^2} \int \Phi_{KLM}^{*l_xl_y}(\Omega_\rho) \times \\ \left(V_{23}^N(|\mathbf{x}_1|) + V_{23}^C(|\mathbf{x}_1|) + V_{31}^C(|\mathbf{x}_2|) + \right. \\ \left. V_{12}^C(|\mathbf{x}_3|) \right) \Phi_{K'LM}^{l'_x l'_y}(\Omega_\rho) d\Omega_\rho \end{aligned} \quad (\text{A15})$$

The system of Eq. (A14) is a system of the linked one dimensional equations which must satisfy the boundary conditions depending on the particular physical situation.

It is easy to show that near $\rho = 0$ regular solutions of (A14) must have the behaviour as ρ^{K+2} . If we omit the nondiagonal terms in the equations, solutions at $\rho \rightarrow \infty$ have the asymptotic behavior as a superposition of the regular and irregular Coulomb functions [17].

-
- [1] J. N. Bahcall, *Neutrino Astrophysics*, Cambridge University Press, 1989.
 - [2] J. N. Bahcall, arXiv:astro-ph/0209080.
 - [3] J. N. Bahcall, and M. H. Pinsonneault, Phys. Rev. Lett. **92**, 121301 (2004).
 - [4] J. N. Bahcall, A. M. Serenelli, and S. Basu, ApJ, **621**, L85 (2005); arXiv:astro-ph/0412440.
 - [5] J. N. Bahcall, A. M. Serenelli, and S. Basu, ApJ Suppl., **165**, 400 (2006).
 - [6] <http://www.sns.ias.edu/~jnb/>
 - [7] J. N. Bahcall and R. M. May, ApJ **155**, 501 (1969).
 - [8] V. B. Belyaev, S. V. Levin, S. L. Yakovlev, J. Phys. **B37**, 1369 (2004).
 - [9] G. Bellini *et al.* (Borexino Collaboration), Phys. Rev. Lett. **108**, 051302 (2012).
 - [10] H. Primakoff, Rev. Mod. Phys. **31**, 802 (1959).
 - [11] B. Povh, K. Rith, C. Scholz, F. Zetsche, *Particles and Nuclei: An Introduction to the Physical Concepts*, Sixth Edition, Springer, 2008.
 - [12] J. M. Blatt, V. F. Weisskopf, *Theoretical nuclear physics*, New York: Springer-Verlag, 1979.
 - [13] <http://www.sns.ias.edu/~jnb/SNdata/Export/BS2005/bs05op.dat>.
 - [14] J. N. Bahcall, A. M. Serenelli, Sarbani Basu, ApJ **621**, L85 (2005); ApJ Suppl. **165**, 400 (2006).
 - [15] E. O. Alt and A. M. Mukhamedzhanov, Phys. Rev. A **47**, 2004 (1993).
 - [16] A. M. Mukhamedzhanov and M. Lieber, Phys. Rev. A **54**, 3078 (1996).
 - [17] R. N. Djibuti and K. V. Shitikova, *Method of hyperspherical functions in atomic and nuclear physics*, Moscow: Energoatomizdat, 1993 (In Russian).
 - [18] M. Fabre de la Ripelle, S. A. Sofianos, R. M. Adam, Annals of Physics **316**, 107 (2005).
 - [19] C. Angulo, *et al.*, Nucl. Phys. A **656**, 3 (1999).